

## Worksheet: Models of Predicate Calculus

---

1. Assume we have a set of function symbols (e.g.,  $f$  or  $\cos$  or  $+$ ), constants (e.g.,  $c$  or  $7$  or  $\text{Bob}$ ), and predicate/relation symbols (e.g.,  $P$  or  $R$  or  $<$ ). A *model* is a specification  $\mathcal{M}$  consisting of:

- A *non-empty* mathematical set  $\mathcal{A}$  (the “individuals”);
- For each constant symbol  $c$ , an element  $c^{\mathcal{M}} \in \mathcal{A}$ ;
- For each function symbol  $f$  with arity  $n$ , a mathematical function  $f^{\mathcal{M}} : \mathcal{A}^n \rightarrow \mathcal{A}$ ;
- For each relation symbol  $R$  with arity  $n$ , a mathematical relation  $R^{\mathcal{M}}$  on the set  $\mathcal{A}^n$  (or, equivalently, a mathematical function  $\mathcal{A}^n \rightarrow \{\text{t}, \text{f}\}$ ).

Suppose we have constants  $c$  and  $d$ , a unary function  $k$  and a binary relation  $R$ . Give three different models for these ingredients.

2. In each of your three models, which of these statements are true? (Use your intuition; for exists and forall, think about all the elements of your set  $\mathcal{A}$ )

- $R(c, k(d))$
- $R(c, d) \vee R(d, c)$
- $\exists x. R(x, x)$
- $\forall x. R(x, x)$
- $\forall x. R(x, x) \vee \neg R(x, x)$

3. Consider the following model  $\mathcal{M}$ : the set  $A$  consists of the people in this classroom, there is one unary predicate  $W$  that is true if the person is awake, and there is one constant  $\text{prof}$  representing the professor.

- If we want to check whether  $A \wedge B$  is true in a model, what do we need to check?

- If we want to check whether  $W(x)$  is true, what else do we need to know?

- If we want to know whether  $\forall x. W(x)$  is true, what do we need to check?

- If we want to know whether  $\exists x. W(x)$  is true in a model, what do we need to check?

4. We say that  $A_1, A_2, \dots \models B$  if every model where all the  $A_i$ 's true also makes  $B$  true. As in predicate logic (although the proof is much harder) we have that

$$A_1, A_2, \dots \models B \quad \text{if and only if} \quad A_1, A_2, \dots \vdash B.$$

Show that the following statements are *not* provable by finding an appropriate model.

- Assumption:  $\forall x. T(x, x)$ . Conclusion:  $\forall x. \forall y. T(x, y)$

- Assumptions:  $D(c)$ , and  $\forall y. C(y) \rightarrow D(y)$ . Conclusion:  $C(c)$ .

- Assumption:  $\forall x. \forall y. \forall z. M(x, y) \wedge M(y, z) \rightarrow M(x, z)$ .  
Conclusion:  $\forall x. \forall y. M(x, y) \vee M(y, x)$ .